



# Anti-nutritional factors: its impacts in animal nutrition

Manju Lata

*Assistant Professor, Department of Animal Nutrition, College of Veterinary and Animal Sciences, GBPUAT, Pantnagar, 263145, Uttarakhand, India*

*Corresponding author: manjundri1@gmail.com*

*Article Received: 24.03.2021*

*Article Published: 26.03.2021*

## ABSTRACT

**Nutrition** is one of the main important factors driving the functioning efficacy, efficiency and evolution of livestock system. Meeting livestock nutritional requirements is extremely important in maintaining acceptable performance of neonatal, growing, finishing and breeding animals. The main challenges nowadays in ruminant production are to reduce feeding cost, improve products quality. The use of unconventional feedstuffs or browse and shrubs may contribute to decrease feeding cost and environmental impact. The problem of feeding such materials is that they contain different levels of anti-nutritional factors (ANFs). Their metabolic products could reduce the availability of one or more nutrients. These ANFs including phenolic compound, phytates, tannins, saponins and oxalate, have his own mechanisms in inhibiting the utilization of such substances. However, many methods aimed to reduce their deleterious effect. These methods could include heating, treatment with PEG ( Polyethylene glycol) or fungi.

**Keywords :** nutrition, anti-nutritional factors, metabolic products, saponins, tannins, phytates

## INTRODUCTION

It is well known that nutrition is one of the main factors driving the functioning efficacy, efficiency and evolution of livestock system. The main challenges nowadays in ruminant production are to reduce feeding cost, improve products quality and diminish the impact of production on environment. The use of unconventional feedstuffs may contribute to decrease feeding cost and environmental impact through reduced methane emissions as well. Not only

that, but shrubby plants can be used to combat desertification, mitigating the effect of drought, allowing soil fixation and enhancing the restoration of the vegetation and the rehabilitation of rangelands. In the meantime, browsing tree foliage plays an important role in ruminant feeding systems in many tropical and Mediterranean environments around the world. They are chiefly as good, cheap sources of nitrogen and energy, which may reduce feeding cost and raise sheep productivity in arid and semi-arid zone. So, to reach that goal, it must exploit all that is available either unconventional feedstuffs or browse and shrubs. However, the problems of feeding such plants or shrubs (Acacia, Leucaena and Atriplex) that they had different levels of anti-nutritional factors.

### ANTI-NUTRITIONAL FACTORS

Anti-nutritional factors (ANFs) are substances that when present in animal feed or water they either by themselves or through their metabolic products reduce the availability of one or more nutrients. Plants contain starch polysaccharides and non starch polysaccharides (NSPs), and some of them are anti-nutritional factors. NSPs contain sugars other than glucose and/or have linkages other than the linkages common in sugar. An example of an NSP is cellulose, that is a chain of glucose molecules, but the molecules have  $\beta$ -(1 $\rightarrow$ 4) bonds between them. The different orientation of the  $\beta$ -bonds (compared to  $\alpha$ -bonds) makes them resistant to digestion by endogenous digestive enzymes of animals. Cell walls in plants contain cellulose and other polysaccharides or non carbohydrate materials such as protein and lignin. Plants contain a mixture of both water-soluble and insoluble NSPs; and the ratio changes with type and stage of maturity of the plant. Cellulose is insoluble in water and is considered fiber. Most NSPs adversely affect digestion in animals, as they affect the viscosity of the material in the digestive tract, and in turn, affects the ability of the digestive enzymes to do their effect, in the end they result in reduced feed efficiency. As the same time the utility of leaves, pods and edible twigs of shrubs and trees is limited as animal fed by the presence of ANFs.

The anti-nutritional factors may **be classified on the basis** of their effects on the nutritional value of feedstuffs, and on the biological response to them in the animal.

### HUISMAN & TOLMAN DIVIDED THE ANTI- NUTRITIONAL FACTORS INTO GROUPS

1. Factors with a depressive effect on protein digestion and on the utilization of protein:  
Example: Protease inhibitors, Tannins and Saponins;
2. Factors that affect mineral utilization: which include phytates.
3. Factors that stimulate the immune system and may cause a damaging hypersensitivity reaction, such as antigenic proteins.

4. Factors with a negative effect on the digestion of carbohydrates:  
Example : amylase inhibitors, phenolic compound and flatulence factors.

Some antinutritional factors are: Non-protein Amino Acids (Mimosine) as in Leucaena, Glycosides (Saponins) as in Acacia, Polyphenolic compounds (Tannins, Lignins) as in all vascular plants, Alkoaois as in Acacia and Oxalate as in Acacia as well.

**Mimosine** : act as an amino acid or make a disruption of the catalytic, transaminases, or may complex with metal such as Zinc. To come over to the mimosine problem when feeding Leucaena, is to restrict to 30% of the green forage with cattle and buffalo, and 50% for goats.

**Saponins** : are heterogenous groups of naturally occurring foam producing steroidal glycosides that occur in a wide range of plants, including oilseeds such as kidney bean, lentil, pea, chickpea, alfalfa, soybean, groundnut and sunflower. They reduce the uptake of certain nutrients including glucose and cholesterol in the gut through intra luminal physicochemical interaction. Hence, they have been reported to have hypocholesterolemic effects. They have distinctive foaming characteristics with white clover and alfalfa; they can cause bloat, hemolysis and inhibit microbial fermentation and synthesis in rumen. However, it has varied biological effects due to structural differences in their sapogenin fractions.

**Tannins** : are water soluble phenolic compounds, they are the most common type found in forage legumes, trees and shrubs. They have the ability to precipitate proteins from aqueous solution. Tree and shrub leaves contain the two different groups (hydrolysable & condensed Tannins (CT)). Tannins have more effect in reducing digestibility than hydrolysable tannins. However, the mechanism effect of tannins came from their ability to form strong H bonds with nutrients resulted in inhibitions of digestive enzymes and rumen microbial activity, and their effect can increased with the increase of the tannins molecular significantly. Concentrations of 2-4% of DM increase N utilization due to increased bypass, Concentrations >7% usually reduce nutrient utilization.

Tannins are present in the NDF and ADF of the tree leaves, which are bound to the cell wall & cell protein and can resulted in decreasing digestibility as they also cause decreased palatability, feed intake, reduced growth rate or loss in weight, poor utilization and decrease iron absorption. But, on the other hand, tannins have some advantages due to their efficiency on animal health, as it had other properties such as, anti -diarrhea, anti- bacterial, anti- oxidant, free radical, scavenging anti- proliferative activity in liver cells.

**Alkaloids** : cause gastrointestinal and neurological disorders. The glyco-alkaloids, solanine and chaconine present in potato and *Solanum* spp are toxic to fungi and humans. Some plant alkaloids are reported to cause infertility.

**Anti vitamin factors** : there are some anti-vitamin factors in some plants, especially leguminous plants.

**Anti-vitamin E**: has also been noted in isolated soya protein, which is suspected to be tocopherol oxidase.

**Anti-metals Phytates** : bind minerals like calcium, iron, magnesium and zinc and make them unavailable. Anemia and other mineral deficiency disorders are common in regions where the diet is primarily a vegetarian.

**Oxalate** : is considered an anti-nutrient because it inhibits calcium absorption and can increase the risk of developing kidney stones.

#### **Methods of reduce the deleterious effect of ANF's in feedstuffs :**

A number of methods have been tried to overcome the deleterious effect of such anti-nutritional factors and tannins is came at the head. These are through making hay, silage with inoculants, using PEG, urea or biological treatment with fungi can be applied to either take off or minimized

and decrease anti-nutritional factors concentration.

#### **Alkali treatment includes polyethylene glycol (PEG):**

which a tannins-binding agent, was shown to be a powerful tool for isolating the effect of tannins on various digestive function. But it may not be economic. Although the incorporation of polyethylene glycol (PEG), which binds with and inactivates tannins, is quite effective, success of its adoption depends on the cost: benefit ratio.

#### **CONCLUSION**

For increasing the utilization of dietary nutrients, reducing environmental contamination and decrease feeding cost, the optimum use of unconventional feedstuffs as well as any local sources (shrubs, browsing tree) has big potential.

#### **REFERENCES**

1. Hegarty. *Toxic amino acids of plant origin*. Effect of poisonous plants on livestock. 1987.
2. Price KR, Johnson IT, Fenwick GR. The chemistry and biological significance of saponins in foods and feeding stuffs. *Crit Rev Food Sci Nutr*. 1987; 26(1):27–135.
3. Min BR, Barry TN, Attwood GT, et al. The effect of condensed tannins on the nutrition and health of ruminants fed fresh lemirate forage: a review. *Anim Feed Sci Technol*. 2003; 106 (1–4):3–19.
4. Kumar R, Singh M. Tannins their adverse role in ruminant nutrition. *J Agric Food Chem*. 1984;32(3):447–453.